

I. Homotopy

Definition 1.1. Two maps $f_i: X \rightarrow Y$, $i = 0, 1$, are called homotopic if there exists a map $F: X \times I \rightarrow Y$ such that $F(x, i) = f_i(x)$ for $i = 0, 1$ and all $x \in X$. Here $I = [0, 1]$. Denote this relation by $f_0 \simeq f_1$.

Proposition 1.2. \simeq is an equivalence relation between maps $X \rightarrow Y$.

Proposition 1.3. \simeq is preserved under composition.

Definition 1.4. $b: I \rightarrow X$ is a path from $b(0)$ to $b(1)$. X is path-connected if

Proposition 1.5. Every path-connected space is connected. There is a compact, connected subset of \mathbb{R}^2 which is not path-connected.

Proposition 1.6. \mathbb{R}^n , S^n , and I^n are path-connected if $n \geq 0$ except for S^0 .

Definition 1.7. $f: X \rightarrow Y$ is a homotopy equivalence if $\exists g: Y \rightarrow X$ such that $gf \simeq id_X$ and $fg \simeq id_Y$. g is called a homotopy inverse to f .

Proposition 1.8. A homotopy inverse to a homotopy equivalence is a homotopy equivalence. The composition of two homotopy equivalences is a homotopy equivalence.

Definition 1.9. X and Y are of the same homotopy type if \exists a homotopy equivalence $f: X \rightarrow Y$.

Proposition 1.10. Being of the same homotopy type is an equivalence relation.

Proposition 1.11. $\exists X$ and Y of the same homotopy type but X and Y are not homeomorphic.

Definition 1.12. X is contractible if X is of the same homotopy type as a one point space P .

Proposition 1.13. \mathbb{R}^n and I^n are contractible if $n \geq 0$.

Proposition 1.14. X is contractible if and only if all maps $X \rightarrow X$ are homotopic.

Definition 1.15. $f: (X,A) \rightarrow (Y,B)$ if $f: X \rightarrow Y$ and $f(A) \subset B$. Define the notion of homotopy between maps of pairs.

Proposition 1.16. \simeq is an equivalence relation.

II. The Fundamental Group

Construction 2.1. If $x \in X$, let $c_x: I \rightarrow X$ be the path defined by $c_x(t) = x$ for all $t \in I$. If b is a path, let $b^{-1}(t) = b(1-t)$ for all $t \in I$. If b and d are paths in X such that $b(1) = d(0)$, define $b * d$ by $b * d(t) = b(2t)$ for $0 \leq t \leq \frac{1}{2}$ and $= d(2t-1)$ for $\frac{1}{2} \leq t \leq 1$. Let $[b]$ denote the homotopy class of all paths in X which are homotopic to b , considered as maps $b: (I, \{0\}, \{1\}) \rightarrow (X, \{b(0)\}, \{b(1)\})$.

Proposition 2.2. If b is a path in X , then $[b * b^{-1}] = [c_{b(0)}]$ and $[b^{-1} * b] = [c_{b(1)}]$.

Proposition 2.3. If $b, d,$ and e are paths in X such that $b(1) = d(0)$ and $d(1) = e(0)$, then $[(b * d) * e] = [b * (d * e)]$.

Proposition 2.4. If $[b] = [b']$ and $[d] = [d']$ and $b(1) = d(0)$, then $[b * d] = [b' * d']$.

Definition 2.5. A path b in X is a loop at $x_0 \in X$ if $b(0) = b(1) = x_0$. $[b]$ denotes the homotopy class of loops homotopic to b .

Theorem 2.6. The homotopy classes of loops at $x_0 \in X$

form a group with the product $[b_1] * [b_2] = [b_1 * b_2]$. This group is called the fundamental group or first homotopy group of X with x_0 as base point and is denoted by $\pi_1(X, x_0)$.

Theorem 2.7. Let $f: (X, x_0) \rightarrow (Y, y_0)$. Define $f_\# : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ by $f_\#([b]) = [fb]$. Then $f_\#$ is a homomorphism.

Theorem 2.8. $(id_X)_\# = id$. If $f: (X, x_0) \rightarrow (Y, y_0)$ and $g: (Y, y_0) \rightarrow (Z, z_0)$, then $g_\#f_\# = (gf)_\#$.

Theorem 2.9. If $f_0 \simeq f_1: (X, x_0) \rightarrow (Y, y_0)$, then $(f_0)_\# = (f_1)_\#$.

Definition 2.10. $f: (X, x_0) \rightarrow (Y, y_0)$ is a homotopy equivalence relative to the base point if $\exists g: (Y, y_0) \rightarrow (X, x_0)$ such that $gf \simeq id_{(X, x_0)}$ and $fg \simeq id_{(Y, y_0)}$.

Proposition 2.11. If f is a homotopy equivalence relative to the base point, then $f_\#$ is an isomorphism.

Corollary 2.12. $\pi_1(\mathbb{R}^n, 0) = \{1\}$ and $\pi_1(I^n, 0) = \{1\}$ if $n \geq 0$.

Theorem 2.13. $\pi_1(X \times Y, (x_0, y_0)) \cong \pi_1(X, x_0) \times \pi_1(Y, y_0)$.

Proposition 2.14. Let $V \subset X$ be the path component of x_0 in X and let $j: (V, x_0) \rightarrow (X, x_0)$ be the inclusion map. Then $j_\#: \pi_1(V, x_0) \rightarrow \pi_1(X, x_0)$ is an isomorphism.

Theorem 2.15. Let $j: I \rightarrow S^1$ be defined by $j(t) = e^{2\pi it}$, and let $s_0 = j(0) = 1$. Define $j^\#: (\text{homotopy classes of maps } (S^1, s_0) \rightarrow (X, x_0)) \rightarrow \pi_1(X, x_0)$ by $j^\#([b]) = [bj]$. Then $j^\#$ is a bijection.

Construction 2.16. Let b be a path in X . Define $b_*: \pi_1(X, b(0)) \rightarrow \pi_1(X, b(1))$ by $b_*(q) = [b^{-1}] * q * [b]$.

Proposition 2.17. b_* is a homomorphism. If d is a path

such that $d(0) = b(1)$, then $d_*b_* = (b * d)_*$. If $[b] = [b']$, then $b_* = b'_*$. $(c_x)_*$ is the identity.

Theorem 2.18. b_* is an isomorphism.

Lemma 2.19. Let $h_0 \simeq h_1: X \rightarrow Y$. Then $\exists b$, a path in Y , such that $b_*h_{0\#} = h_{1\#}$.

Proposition 2.20. Let $f: X \rightarrow Y$ be a homotopy equivalence. Then $f_{\#}: \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))$ is an isomorphism. (Compare 2.11.)

Definition 2.21. X is simply connected (or 1-connected) if X is path connected and $\pi_1(X, x_0) = \{1\}$ for some $x_0 \in X$.

Proposition 2.22. If X is simply connected, then $\pi_1(X, x_0) = \{1\}$ for all $x_0 \in X$.

Proposition 2.23. Every contractible space is simply connected.

Theorem 2.24. Let b be a path from x_0 to x_1 . Then $b_*: \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$ is independent of the choice of b if and only if $\pi_1(X, x_0)$ is abelian.

Definition 2.25. An H-space is a pair (H, e) , where H is a space, $e \in H$, and we are given a map $m: H \times H \rightarrow H$ such that $m(\cdot, e) \simeq m(e, \cdot) \simeq \text{id}_{(H, e)}$.

Example 2.26. A topological group is an H-space.

Theorem 2.27. Let (H, e) be an H-space. Then $\pi_1(H, e)$ is abelian.

III. Covering Spaces

Definition 3.1. A space X is called locally pathwise connected if for every $x \in X$ and every open set U with $x \in U$, there exists a pathwise connected open set V such that $x \in V$

C U.

Definition 3.2. Let $f: Y \rightarrow X$ and let U be an open set of X . Then U is evenly covered by f if there is an indexing set J and for every $j \in J$, an open set $U_j \subset Y$ such that

- (i) $\bigcup_{j \in J} U_j = f^{-1}(U)$,
- (ii) if $j \neq k$, $U_j \cap U_k = \emptyset$,
- (iii) $f|_{U_j}: U_j \rightarrow U$ is a homeomorphism for $j \in J$.

Definition 3.3. A map $\pi: X^{\wedge} \rightarrow X$ is a covering of X if

- (i) π is onto,
- (ii) X^{\wedge} is pathwise connected and locally pathwise connected,
- (iii) for every $x \in X$, \exists an open U which is evenly covered by π and $x \in U$.

Proposition 3.4. The following are covering spaces.

- (i) $\text{id}_X: X \rightarrow X$, if X is pathwise connected and locally pathwise connected,
- (ii) $\pi: \mathbb{R}^1 \rightarrow S^1$ given by $\pi(r) = e^{2\pi i r}$,
- (iii) $\pi_n: S^1 \rightarrow S^1$ given by $\pi_n(e^{i r}) = e^{i n r}$, for $n \neq 0$ an integer,
- (iv) $\pi: S^n \rightarrow \mathbb{R}P^n = \{\text{pairs of antipodal points of } S^n\}$ with the quotient topology ($\mathbb{R}P^n$ is called real n -dimensional projective space).

Notation. l.c. stands for pathwise connected and locally pathwise connected.

Proposition 3.5. Let $U \subset X$ be evenly covered by $f: Y \rightarrow X$ and let Y be l.c. Then U is locally pathwise connected.

Proposition 3.6. Let $\pi: X^{\wedge} \rightarrow X$ be a covering. Then X is l.c.

Proposition 3.7. Let $U \subset X$ be evenly covered by $f: Y \rightarrow X$. Let Z be connected and let $g: Z \rightarrow U$. Let $y \in Y$, $z \in Z$ be such that $f(y) = g(z)$. Then $\exists!$ $g': Z \rightarrow Y$ such that $g'(z) = y$ and $fg' = g$.

Proposition 3.8. Let $\pi: X^\wedge \rightarrow X$ be a covering. Let b be a path in X and let $x^\wedge \in X^\wedge$ be such that $\pi(x^\wedge) = b(0)$. Then $\exists!$ path b^\wedge in X^\wedge such that $x^\wedge = b^\wedge(0)$ and $\pi b^\wedge = b$.

Proposition 3.9. Let $\pi: X^\wedge \rightarrow X$ be a covering. Let Z be pathwise connected and let $g_i: Z \rightarrow X^\wedge$ be maps such that $\pi g_0 = \pi g_1$ and $g_0(z) = g_1(z)$ for some $z \in Z$. Then $g_0 = g_1$.

Proposition 3.10. Let $\pi: X^\wedge \rightarrow X$ be a covering, let $g: I^2 \rightarrow X$ be a map, and let $x^\wedge \in X^\wedge$ be such that $\pi(x^\wedge) = g(0,0)$. Then $\exists!$ $g^\wedge: I^2 \rightarrow X^\wedge$ such that $g^\wedge(0,0) = x^\wedge$ and $\pi g^\wedge = g$.

Proposition 3.11. Let $\pi: X^\wedge \rightarrow X$ be a covering. Then $\pi_\# : \pi_1(X^\wedge, x^\wedge) \rightarrow \pi_1(X, \pi(x^\wedge))$ is one to one.

IV: Classification of Coverings with Base Point

Definition 4.1. $\pi: (X^\wedge, x^\wedge) \rightarrow (X, x)$ is a covering with base point if $\pi: X^\wedge \rightarrow X$ is a covering.

Definition 4.2. Two coverings $\pi_i: (X^\wedge_i, x^\wedge_i) \rightarrow (X, x)$ are homeomorphic if \exists a homeomorphism $h: (X^\wedge_1, x^\wedge_1) \rightarrow (X^\wedge_2, x^\wedge_2)$ such that $\pi_2 h = \pi_1$.

Proposition 4.3. Being homeomorphic is an equivalence relation on the set of coverings with base point.

Proposition 4.4. Let π_1 and π_2 be homeomorphic coverings. Then $(\pi_1)_\#(\pi_1(X^\wedge_1, x^\wedge_1)) = (\pi_2)_\#(\pi_1(X^\wedge_2, x^\wedge_2)) \subset \pi_1(X, x)$.

Proposition 4.5. Let π be a covering and let $g: (Z, z) \rightarrow (X, x)$ be a map such that $\text{Im } g_{\#} \subset \text{Im } \pi_{\#}$. Let b and d be paths in Z such that $b(0) = d(0) = z$ and $b(1) = d(1) = z_1$. Let b^{\wedge} and d^{\wedge} be the unique paths in X^{\wedge} such that $\pi b^{\wedge} = gb$, $\pi d^{\wedge} = gd$, and $b^{\wedge}(0) = d^{\wedge}(0) = x^{\wedge}$. Then $b^{\wedge}(1) = d^{\wedge}(1)$.

Proposition 4.6. Let π be a covering, let Z be l.c., and let $g: (Z, z) \rightarrow (X, x)$ be a map such that $\text{Im } g_{\#} \subset \text{Im } \pi_{\#}$. Then $\exists!$ $g^{\wedge}: (Z, z) \rightarrow (X^{\wedge}, x^{\wedge})$ such that $\pi g^{\wedge} = g$.

Proposition 4.7. Two coverings π_1 and π_2 are homeomorphic if and only if $\text{Im } (\pi_1)_{\#} = \text{Im } (\pi_2)_{\#}$.

Definition 4.8. X is semi-locally 1-connected if for every $x \in X$ there is an open set U , $x \in U$, such that $\pi_1(U, x) \rightarrow \pi_1(X, x)$ is the trivial homomorphism.

Definition 4.9. Let $G \subset \pi_1(X, x)$ be a subgroup. Two paths b and d are G-homotopic if $b(0) = d(0) = x$, $b(1) = d(1)$, and $[b * d^{-1}] \in G$.

Proposition 4.10. G-homotopy is an equivalence relation on paths in X which start at x .

Construction 4.11. Let X_G denote the set of G-homotopy classes of paths in X which start at x . Let $\pi: X_G \rightarrow X$ be defined by $\pi(\langle b \rangle) = b(1)$. Let $V \subset X$ be open and path connected. Let b be a path in X such that $b(0) = x$ and $b(1) \in V$. Define $U(b, V) \subset X_G$ to consist of the G-homotopy classes of paths of the form $b * d$ where $d(0) = b(1)$ and $d(t) \in V$. Take the $U(b, V)$ as a subbase for the open sets of X_G .

Theorem 4.12. Let X be l.c. and semi-locally 1-connected. Then the homeomorphism classes of coverings of (X, x) are in one to one correspondence (i. e. \exists a bijection)

with the subgroups of $\pi_1(X, x)$.

V. Classification of Coverings and Their Translations.

Definition 5.1. Two coverings $\pi_i: X^{\wedge}_i \rightarrow X$, $i = 1, 2$, are homeomorphic if there is a homeomorphism $h: X^{\wedge}_1 \rightarrow X^{\wedge}_2$ such that $\pi_2 h = \pi_1$.

Proposition 5.2. Being homeomorphic is an equivalence relation on the set of coverings of X .

Definition 5.3. Two subgroups H, K , of a group G are conjugate if there is a $g \in G$ such that $K = g^{-1}Hg$.

Proposition 5.4. Being conjugate is an equivalence relation on the set of subgroups of G .

Proposition 5.5. Let π be a covering and let $x^{\wedge}_i \in X^{\wedge}$ be such that $\pi(x^{\wedge}_i) = x$, $i = 1, 2$. Then $\pi_{\#}(\pi_1(X^{\wedge}, x^{\wedge}_1))$ and $\pi_{\#}(\pi_1(X^{\wedge}, x^{\wedge}_2))$ are conjugate in $\pi_1(X, x)$.

Proposition 5.6. Let π be a covering, let $x^{\wedge} \in X^{\wedge}$, and let $G \subset \pi_1(X, \pi(x^{\wedge}))$ be conjugate with $\text{Im } \pi_{\#}$. Then $\exists x^{\wedge}_1 \in X^{\wedge}$ such that $\pi_{\#}(\pi_1(X^{\wedge}, x^{\wedge}_1)) = G$.

Theorem 5.7. Let π_i be coverings and let $x^{\wedge}_i \in X^{\wedge}_i$ be such that $\pi_1(x^{\wedge}_1) = \pi_2(x^{\wedge}_2)$. Then π_1 and π_2 are homeomorphic if and only if $\text{Im } (\pi_1)_{\#}$ and $\text{Im } (\pi_2)_{\#}$ are conjugate in $\pi_1(X, \pi_1(x^{\wedge}_1))$.

Theorem 5.8. The homeomorphism classes of coverings of a l.c. and semi-locally 1-connected space X are in one to one correspondence with the conjugacy classes of subgroups of $\pi_1(X, x)$.

Definition 5.9. A translation of a covering π is a homeomorphism $h: X^{\wedge} \rightarrow X^{\wedge}$ such that $\pi h = \pi$. Let $T(\pi)$ denote

the set of translations of π .

Proposition 5.10. $T(\pi)$ is a group under composition of translations.

Definition 5.11. Let $H \subset G$ be a subgroup of G . The elements $g \in G$ such that $g^{-1}Hg = H$ form a subgroup $N(H)$, the normalizer of H in G .

Proposition 5.12. H is normal in $N(H)$.

Construction 5.13. Let π be a covering and let $H = \pi_{\#}(\pi_1(X^{\wedge}, x^{\wedge}))$. Define a function $t: N(H) \rightarrow T(\pi)$ as follows. Let $q \in N(H)$. Choose a path b^{\wedge} in X^{\wedge} such that $b^{\wedge}(0) = x^{\wedge}$ and $[\pi b^{\wedge}] = q$. $t(q)$ is the unique translation such that $t(q)(x^{\wedge}) = b^{\wedge}(1)$.

Theorem 5.14. t is well defined, t is a homomorphism onto $T(\pi)$, and $\text{Ker } t = H$.

Corollary 5.15. $N(H)/H$ is isomorphic to $T(\pi)$.

VI. Some Fundamental Groups.

Definition 6.1. A covering π is called universal if X^{\wedge} is simply connected.

Corollary 6.2. Let π be a universal covering. Then $\pi_1(X, x)$ is isomorphic to $T(\pi)$.

Proposition 6.3. Example 3.4 (ii) is a universal covering.

Theorem 6.4. $\pi_1(S^1, s)$ is isomorphic to the additive group of integers, \mathbb{Z} , for any $s \in S^1$.

Example 6.5. Describe the covering space corresponding to each subgroup of $\pi_1(S^1, s)$.

Definition 6.6. $T^n = S^1 \times \dots \times S^1$, n times, is the

n-dimensional torus.

Theorem 6.7. $\pi_1(T^n, t) \cong \mathbb{Z} \oplus \dots \oplus \mathbb{Z}$, n times.

Theorem 6.8. Let X be l.c. and semi-locally 1-connected, let $A, B \subset X$ be open subsets which are l.c. and simply-connected, and let $A \cap B$ be pathwise connected and $A \cup B = X$. Then X is simply-connected.

Theorem 6.9. S^n is simply-connected for all $n > 1$.

Theorem 6.10. Example 3.4 (iv) is a universal covering and hence $\pi_1(\mathbb{R}P^n, p)$ is isomorphic to $\mathbb{Z}/2\mathbb{Z}$ if $n > 1$.

VII. Higher Homotopy Groups.

Definition 7.1. Y^X denotes the function space of continuous maps $f: X \rightarrow Y$ with the compact-open topology. If $B \subset Y$ and $A \subset X$, then $(Y, B)^{(X, A)} \subset Y^X$ is the subspace of functions such that $f(A) \subset B$.

Theorem 7.2. Define $\alpha: Z^X \times Y \rightarrow (Z^Y)^X$ by $[\alpha(f)](x)(y) = f(x, y)$. If Y is locally compact and Hausdorff, then α is a bijection.

Definition 7.3. $(X, x_0)^{(1, 0)}$ is the space of paths on X starting at x_0 . $(X, x_0)^{(1, (0, 1))}$ is the space of loops on X at x_0 and denoted by $\Omega(X, x_0)$. Define $\Omega^n(X, x_0) = \Omega(\Omega^{n-1}(X, x_0), x_{n-1})$, where x_{n-1} is the constant loop at x_{n-2} .

Definition 7.4. $\pi_n(X, x_0) = \pi_1(\Omega^{n-1}(X, x_0), x_{n-1})$.

Theorem 7.5. $\Omega(X, x_0)$ is an H-space.

Corollary 7.6. $\pi_n(X, x_0)$ is abelian if $n > 1$.

Construction 7.7. Let $f: (X, x_0) \rightarrow (Y, y_0)$. Define $f_1: \Omega(X, x_0) \rightarrow \Omega(Y, y_0)$ by $f_1(b) = fb$. Define $f_n: (\Omega^n(X, x_0), x_n) \rightarrow (\Omega^n(Y, y_0), y_n)$ by induction. Define $f_{\#}: \pi_n(X, x_0) \rightarrow$

$\pi_n(Y, y_0)$ by $(f_{n-1})_{\#}: \pi_1(\Omega^{n-1}(X, x_0), x_{n-1}) \rightarrow \pi_1(\Omega^{n-1}(Y, y_0), y_{n-1})$.

Theorem 7.8. $\text{id}_{\#}$ is the identity. If $f: (X, x_0) \rightarrow (Y, y_0)$ and $g: (Y, y_0) \rightarrow (Z, z_0)$, then $g_{\#}f_{\#} = (gf)_{\#}$.

Proposition 7.9. If $f \simeq g: (X, x_0) \rightarrow (Y, y_0)$, then $f_{\#} = g_{\#}$.

Corollary 7.10. If (X, x_0) is contractible, then $\pi_n(X, x_0) = 0$ for $n \geq 1$.

Theorem 7.11. $\pi_n(X, x_0) \cong \pi_{n-1}(\Omega(X, x_0), x_1)$.

Definition 7.12. Let $I' \subset I^n$ denote the subspace of all n -tuples for which at least 1 coordinate is 0 or 1. $\pi(I^n, I'; X, x_0)$ denotes the set of homotopy classes of maps from (I^n, I') to (X, x_0) .

Theorem 7.13. There is a natural 1-1 correspondence between $\pi_n(X, x_0)$ and $\pi(I^n, I'; X, x_0)$.

VIII. Fibre Spaces.

Definition 8.1. Let $p: E \rightarrow B$. p is a fibre map if for every l.c. space X the outer triangle in the diagram below can be filled in by the dotted arrow so that the triangles commute

$$\begin{array}{ccccc}
 & & & E & \\
 & & & | & \\
 & & & \downarrow & \\
 & & G & & \\
 g & & & & \\
 X & \xrightarrow{\quad} & X \times I & \xrightarrow{\quad} & B \\
 i_0 & & & H &
 \end{array}$$

where $i_0(x) = (x, 0)$. (E, p, B) is called a fibre space, E is the total space, B the base space, and p the projection. If B has a base point b_0 , then $F = p^{-1}(b_0)$ is the fibre.

Proposition 8.2. If B is pathwise connected and $E \neq \emptyset$, then p is onto.

Proposition 8.3. Let $E = F \times B$ and let p be the projection onto the second factor. Then (E, p, B) is a fibre space.

Proposition 8.4. A covering space (X^{\wedge}, π, X) is a fibre space.

Theorem 8.5. Let $A, B \subset Y$. Let $E(Y; A, B) \subset Y^I$ be the subspace of those maps such that $f(0) \in A$ and $f(1) \in B$. Define $p: E(Y; A, B) \rightarrow A \times B$ by $p(f) = (f(0), f(1))$. Then p is a fibre map.

Corollary 8.6. $p: (X, x_0)^{(I, 0)} \rightarrow X$ by $p(f) = f(1)$ is a fibre space with fibre $\Omega(X, x_0)$.

IX. Exact Sequences.

Definition 9.1. A sequence $\dots \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow \dots$ of abelian groups and homomorphisms is exact if $\text{Im } f = \text{Ker } g$.

Proposition 9.2. $0 \rightarrow A \rightarrow 0$ is exact if and only if $A = 0$.

Proposition 9.3. $0 \rightarrow A \xrightarrow{f} B \rightarrow 0$ is exact if and only if f is an isomorphism.

Proposition 9.4. $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ is exact if and only if $g^{\wedge}: B/f(A) \rightarrow C$ is an isomorphism and f is injective.

"Five Lemma" 9.5. Let

$$\begin{array}{ccccccccc}
 A & \rightarrow & B & \rightarrow & C & \rightarrow & D & \rightarrow & E \\
 & & f & & g & & h & & i \\
 | & & | & & | & & | & & | \\
 |P_1 & & |P_2 & & |P_3 & & |P_4 & & |P_5
 \end{array}$$

$$\begin{array}{ccccccc}
 & \downarrow & & \downarrow & & \downarrow & \\
 A' & \rightarrow & B' & \rightarrow & C' & \rightarrow & D' & \rightarrow & E' \\
 & & f' & & g' & & h' & & i'
 \end{array}$$

be a commutative diagram with exact rows. If p_1 , p_2 , p_4 , and p_5 are isomorphisms, so is p_3 .

X. Relative Homotopy Groups and Exact Sequences.

Definition 10.1. Let $I^{n-1} \subset I^n$ be the subspace where $t_n = 0$. Let $J^{n-1} \subset I^n$ be the closure of $I^n - I^{n-1}$. Let $x_0 \in A \subset X$. For $n \geq 2$, define $\pi_n(X, A)$ to be the set of homotopy classes of maps from $(I^n, I^{n-1}, J^{n-1}) \rightarrow (X, A, x_0)$.

Proposition 10.2. Let $Z \subset X^I$ be defined by $Z = \{f \mid f(0) = x_0, f(1) \in A\}$. Then $\pi_n(X, A) \cong \pi_{n-1}(Z, c)$.

Corollary 10.3. $\pi_n(X, A)$ is abelian for $n > 2$.

Definition 10.4. Let $f: (X, A, x_0) \rightarrow (Y, B, y_0)$. Define $f_\#: \pi_n(X, A, x_0) \rightarrow \pi_n(Y, B, y_0)$ in the usual way.

Theorem 10.5. $f_\#$ is a homomorphism, $(\text{id})_\# = \text{id}$, and $g_\# f_\# = (gf)_\#$.

Definition 10.6. $\partial: \pi_n(X, A) \rightarrow \pi_{n-1}(A, x_0)$ is defined by $\partial([f]) = [f|_{I^{n-1}}]$.

Theorem 10.7. Let $i: (A, x_0) \rightarrow (X, x_0)$, $j: (X, x_0) \rightarrow (X, A)$.

Then

$\dots \rightarrow \pi_n(A, x_0) \xrightarrow{\partial} \pi_n(X, x_0) \xrightarrow{i_\#} \pi_n(X, A) \xrightarrow{j_\#} \pi_{n-1}(A, x_0) \xrightarrow{\partial} \dots$
is exact. (This is called the exact sequence of a pair.)

Theorem 10.8. Let (E, p, B) be a fibre space, $b_0 \in B$, $p^\wedge: (E, F) \rightarrow (B, b_0)$. Then $p^\wedge_\#: \pi_n(E, F) \rightarrow \pi_n(B, b_0)$ is an isomorphism for $n \geq 2$.

Theorem 10.9. Let (E, p, B) be a fibre space, $b_0 \in B$, $e_0 \in$

F. Then there is an exact sequence

$$\dots \xrightarrow{\partial^\wedge} \pi_n(F, e_0) \xrightarrow{p_\#} \pi_n(E, e_0) \xrightarrow{\partial^\wedge} \pi_n(B, b_0) \xrightarrow{p_\#} \pi_{n-1}(F, e_0) \xrightarrow{\partial^\wedge} \dots$$

Here $\partial^\wedge = \partial(p^\wedge_\#)^{-1}$. (This is called the exact sequence of a fibre space.)

Theorem 10.10. Let (X^\wedge, π, X) be a covering space. Then

$\pi_\#: \pi_n(X^\wedge, x^\wedge) \rightarrow \pi_n(X, \pi(x^\wedge))$ is an isomorphism for $n \geq 2$.

Corollary 10.11. $\pi_n(S^1, x_0) = 0$ if $n > 1$.

Corollary 10.12. $\pi_n(S^n, x_0) \cong \pi_n(\mathbb{R}P^n, (x_0, -x_0))$, if $n > 1$.

11. Polyhedra

Definition 11.1. A simplicial complex K (of dimension at most 2) is a collection of vertices $\{V_i\}$, a collection of edges $\{V_i, V_j\}$, and a collection of triangles $\{V_i, V_j, V_k\}$ satisfying the following condition: given a set in the collection, then a non-empty subset of it is in the collection.

(Intuitive) Definition 11.2. A topological space X is a polyhedron if it can be broken up into vertices, edges, and triangles as in definition 11.1. More precisely, if X is homeomorphic to $|K|$, the geometric realization of K .

Proposition 11.3. I^n , S^n , and \mathbb{R}^n are polyhedra (if $n \leq 2$).

Definition 11.4. Given a simplicial complex K . An edge path in K is a sequence of vertices of K , $w = V_0 \dots V_k$, such that each successive pair of vertices is an edge or repeats itself. If $w' = V_k \dots V_1$, then $w * w' = V_0 \dots V_1$, the product of edge-paths. $w^{-1} = V_k \dots V_0$ is the inverse edge-path to w .

Definition 11.5. We define an equivalence relation on

edge-paths with the same beginning and end points as follows.

If $w = v_0 \dots v_i v_j \dots v_k$, then $w \simeq w' = v_0 \dots v_i \dots v_k$. If

v_i, v_j, v_l is a ^{Simplex} triangle in K , then $w = v_0 \dots v_i v_l \dots v_k \simeq w' = v_0 \dots v_i v_j v_l \dots v_k$. Let $[w]$ denote the equivalence class of w .

Proposition 11.6. Let $\pi(K, v)$ denote the set of equivalence classes of edge-paths in K which start and end at v . Then $\pi(K, v)$ is a group under the operation $[w] * [w'] = [w * w']$. This is called the edge-path group of K .

Theorem 11.7. Let X be a polyhedron with corresponding simplicial complex K . Then $\pi(K, v) \cong \pi_1(X, v)$.

Definition 11.8. K is connected if $|K|$ is path-connected. K is simply connected if $|K|$ is simply connected.

Proposition 11.9. Given K . Then there exists a simply ^{connected} subcomplex L (having only vertices and edges) of K which contains all the vertices of K . Such a subcomplex is a maximal tree.

Definition 11.10. Given K and a subcomplex L . Let $G = G(K, L)$ be the group generated by g_{ij} , one generator for each edge v_i, v_j in K subject to the relations $g_{ij} = 1$ if v_i, v_j is in L and $g_{ij}g_{jk} = g_{ik}$ if v_i, v_j, v_k is in K .

Theorem 11.11. If L is a maximal tree, then $\pi(K, v) \cong G(K, L)$. ^{simply connected}

Corollary 11.12. If K is a finite simplicial complex, then $\pi_1(|K|, v)$ is finitely generated and finitely presented.

Corollary 11.13. If K has no triangles, then $\pi_1(|K|, v)$ is free.

Proposition 11.14. π_1 of a figure 8 is free on two

generators.

Theorem 11.15. A free group on two generators contains a free group on n generators.