

$$\mathbf{a} = (a_0, a_1, a_2, \dots)$$

ordinary generating function of

a :

$$a_0 + a_1x + a_2x^2 + \dots = \sum_{n \geq 0} a_n x^n$$

exponential generating function

of a :

$$a_0 + a_1 \frac{x}{1!} + a_2 \frac{x^2}{2!} + \dots = \sum_{n \geq 0} a_n \frac{x^n}{n!}$$

Many others, not as important.

What is the point?

“Natural” algebraic operations on generating functions have combinatorial significance, so we can transform combinatorics into algebra (and vice versa).

Notation:

$$\mathbb{N} = \{0, 1, 2, \dots\}$$

$$\mathbb{P} = \{1, 2, 3, \dots\}$$

$$[n] = \{1, 2, \dots, n\}$$

$$[x^n] \sum a_k x^k = a_n.$$

Some operations:

$$\sum a_n x^n + \sum b_n x^n = \sum (a_n + b_n) x^n$$

$$\left(\sum a_n x^n \right) \left(\sum b_n x^n \right) = \sum c_n x^n,$$

where $c_n = \sum_{k=0}^n a_k b_{n-k}$.

$$\left(\sum a_n \frac{x^n}{n!} \right) \left(\sum b_n \frac{x^n}{n!} \right) = \sum c_n \frac{x^n}{n!},$$

where $c_n = \sum_{k=0}^n \binom{n}{k} a_k b_{n-k}$.

Define

$$G(x) = \mathbf{1/F(x)}$$

if $F(x)G(x) = 1$ (exists if and only if $F(0) \neq 0$). E.g,

$$\frac{1}{1 - ax} = 1 + ax + a^2x^2 + \dots$$

Let

$$F(x) = \sum_{n \geq 0} a_n x^n, \quad G(x) = \sum_{n \geq 1} b_n x^n$$

(so $G(0) = 0$). Define the **composition** $F(G(x))$ by

$$F(G(x)) = \sum_{n \geq 0} a_n G(x)^n.$$

Makes sense **formally** since computing $[x^n]F(G(x))$ involves only a **finite** sum.

Examples. Let $G(0) = 0$. Then

$$e^{G(x)} = \sum_{n \geq 0} \frac{G(x)^n}{n!}$$

$$-\log(1 - G(x)) = \sum_{n \geq 1} \frac{G(x)^n}{n}.$$

Lifting principle: All “familiar” formulas for convergent power series continue to hold whenever they make sense formally. E.g., if $G(0) = 0$ then

$$\begin{aligned} \log(e^{G(x)}) &= G(x) \\ e^{\log(1+G(x))} &= 1 + G(x). \end{aligned}$$

Sets. Let $n \in \mathbb{N}$ and

$$F_n(\mathbf{x}) = \sum_{T \subseteq [n]} \prod_{i \in T} x_i,$$

a “list” of all subsets of $[n]$. E.g.,

$$F_2(\mathbf{x}) = 1 + x_1 + x_2 + x_1x_2.$$

Since for each $i \in S$ either $i \in T$ or $i \notin T$, we have

$$F_n(\mathbf{x}) = (1 + x_1)(1 + x_2) \cdots (1 + x_n).$$

Define

$$\binom{n}{k} = \#\{T \subseteq S : \#T = k\}.$$

Put each $x_i = x$ to get

$$(1 + x)^n = \sum_{k \geq 0} \binom{n}{k} x^k.$$

Illustrates technique of “late specialization.”

Multisets. A **multiset** M on a set S is a set **with repeated elements** from S . E.g,

$$\{1, 1, 1, 2, 4, 4, 4, 7, 7\} = \{1^3, 2, 4^3, 7^2\}$$

is a multiset on $[10]$. Let

$$\nu_M(i) = \# \text{ } i\text{'s in } M.$$

Let

$$G_n(\mathbf{x}) = \sum_{M \text{ on } [n]} \prod_{i=1}^n x_i^{\nu_M(i)},$$

a “list” of all multisets on $[n]$. E.g.,

$$\begin{aligned} G_2(\mathbf{x}) &= 1 + x_1 + x_2 + x_1^2 + x_1x_2 + x_2^2 + \cdots \\ &= (1 + x_1 + x_1^2 + \cdots)(1 + x_2 + x_2^2 + \cdots) \\ &= \frac{1}{(1 - x_1)(1 - x_2)}. \end{aligned}$$

In general,

$$G_n(\mathbf{x}) = \frac{1}{(1 - x_1)(1 - x_2) \cdots (1 - x_n)}.$$

Let $\binom{n}{k}$ denote the number of k -element multisets on $[n]$. E.g., $\binom{3}{2} = 6$:

$$11 \quad 22 \quad 33 \quad 12 \quad 13 \quad 23$$

Put $x_i = x$ to get

$$\begin{aligned} \sum_{k \geq 0} \binom{n}{k} x^k &= \frac{1}{(1 - x)^n} \\ &= \sum_{k \geq 0} \binom{-n}{k} (-x)^k, \end{aligned}$$

where

$$\binom{t}{k} = \frac{t(t - 1) \cdots (t - k + 1)}{k!}.$$

$$\begin{aligned} \sum_{k \geq 0} \binom{\binom{n}{k}}{k} x^k &= \frac{1}{(1-x)^n} \\ &= \sum_{k \geq 0} \binom{-n}{k} (-x)^k, \end{aligned}$$

Hence

$$\binom{\binom{n}{k}}{k} = (-1)^k \binom{-n}{k} = \binom{n+k-1}{k}$$

(example of **reciprocity**).

Combinatorial or **bijjective** proof
that

$$\left(\binom{n}{k}\right) = \binom{n+k-1}{k}$$

Let

$$1 \leq a_1 \leq a_2 \leq \cdots \leq a_k \leq n$$

be a k -multiset on $[n]$. Let $b_i = a_i + i - 1$. Then

$$1 \leq b_1 < b_2 < \cdots < b_k \leq n + k - 1,$$

and conversely (i.e., $a_i = b_i - i + 1$).

Thus

$$\left(\binom{n}{k}\right) = \binom{n+k-1}{k}.$$

RATIONAL GENERATING FUNCTIONS

A generating function $F(x) = \sum a_n x^n$ is **rational** if there are polynomials $P(x), Q(x)$ such that

$$F(x) = \frac{P(x)}{Q(x)},$$

i.e., $F(x)Q(x) = P(x)$. Can assume $Q(0) = 1$.

E.g.,

$$\sum_{n \geq 0} a^n x^n = \frac{1}{1 - ax}.$$

More generally,

$$\begin{aligned} \frac{1}{(1 - ax)^d} &= \sum_{n \geq 0} \binom{-d}{n} (-ax)^n \\ &= \sum_{n \geq 0} \binom{n + d - 1}{d - 1} a^n x^n. \end{aligned}$$

Note: $\binom{n+d-1}{d-1}$ is a polynomial in n of degree $d - 1$.

Fundamental theorem on rational generating functions.

Fix $\alpha_1, \dots, \alpha_d \in \mathbb{C}$, $\alpha_d \neq 0$.

Let $f : \mathbb{N} \rightarrow \mathbb{C}$. TFAE:

- $$\sum_{n \geq 0} f(n)x^n = P(x)/Q(x),$$

where $Q(x) = 1 + \alpha_1x + \dots + \alpha_dx^d$,

$$P(x) \in \mathbb{C}[x],$$

$$\deg(P) < \deg(Q) = d.$$

- For all $n \geq 0$,

$$f(n+d) + \alpha_1f(n+d-1) + \dots + \alpha_df(n) = 0$$

(linear recurrence with constant coefficients).

- For all $n \geq 0$,

$$f(n) = \sum_{i=1}^k P_i(n) \gamma_i^n,$$

where

$$1 + \alpha_1 x + \cdots + \alpha_d x^d = \prod_{i=1}^k (1 - \gamma_i x)^{d_i},$$

the γ_i 's are distinct, and

$$P_i(n) \in \mathbb{C}[n], \quad \deg(P_i) < d_i.$$

Idea of proof. Use partial fractions to write $P(x)/Q(x)$ as linear combination of terms $(1 - \gamma_i x)^e$, $e < d_i$.

What if $\deg P \geq \deg Q$? Then write (uniquely)

$$\frac{P(x)}{Q(x)} = L(x) + \frac{R(x)}{Q(x)},$$

where $L(x), R(x) \in \mathbb{C}[x]$ and


$$\deg R(x) < \deg Q(x).$$

Thus $L(x)$ records the “exceptional values” (finitely many) where the fundamental theorem fails.

Example (the **transfer-matrix method**).

Let $f(n)$ be the number of sequences $a_1 \cdots a_n$, $a_i = 1, 2, 3$, with no $a_i a_{i+1} = 11$ or 23 . Thus

$f(n) = \#$ paths of length $n - 1$ in:



Adjacency matrix: $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$.

Thus $\left(A^k\right)_{ij}$ is the number of paths of length k from i to j , so

$$f(n) = \sum_{i,j=1}^3 \left(A^{n-1}\right)_{ij}.$$

$$\begin{aligned} \Rightarrow \sum_{n \geq 0} f(n+1)x^n &= \sum_{i,j=1}^3 \left(\sum_{n \geq 0} A^n x^n \right)_{ij} \\ &= \sum_{i,j=1}^3 (I - xA)^{-1}_{ij}. \end{aligned}$$

Let $(B : j, i)$ denote the matrix B with row j and column i removed. Then

$$B_{ij}^{-1} = (-1)^{i+j} \frac{\det(B; j, i)}{\det(B)},$$

so

$$\begin{aligned} \sum_{n \geq 0} f(n+1)x^n &= \frac{\sum (-1)^{i+j} \det(I - xA : j, i)}{\det(I - xA)} \\ &= \frac{\mathbf{3 + x - x^2}}{\mathbf{1 - 2x - x^2 + x^3}}. \end{aligned}$$

EXPONENTIAL GENERATING FUNCTIONS

Given $f : \mathbb{N} \rightarrow \mathbb{C}$, write

$$E_f(x) = \sum_{n \geq 0} f(n) \frac{x^n}{n!}.$$

Proposition. Given $f, g : \mathbb{N} \rightarrow \mathbb{C}$,
define $h : \mathbb{N} \rightarrow \mathbb{C}$ by

$$h(\#X) = \sum_{(S,T)} f(\#S)g(\#T),$$

where $\#X < \infty$ and $S, T \subseteq X$ such
that

$$S \cup T = X, \quad S \cap T = \emptyset.$$

Then

$$E_f(x)E_g(x) = E_h(x).$$

Proof. Let $\#X = n$. There are $\binom{n}{k}$ pairs (S, T) with $\#S = k$ and $\#T = n - k$. Hence

$$\begin{aligned} h(n) &= \sum_{k=0}^n \binom{n}{k} f(k)g(n-k) \\ &= \left[\frac{x^n}{n!} \right] E_f(x)E_g(x). \quad \square \end{aligned}$$

Example. Find the number $h(n)$ of ways to let $[n] = S \cup T$ with $S \cap T = \emptyset$, choose a subset of S , and choose an element of T . Here $f(n) = 2^n$ and $g(n) = n$. Thus

$$E_f(x) = \sum_{n \geq 0} 2^n \frac{x^n}{n!} = e^{2x}$$

$$E_g(x) = \sum_{n \geq 0} n \frac{x^n}{n!} = xe^x$$

$$\begin{aligned} \Rightarrow E_h(x) &= xe^{3x} \\ &= \sum_{n \geq 0} n 3^{n-1} \frac{x^n}{n!}, \end{aligned}$$

whence $h(n) = n3^{n-1}$.

Iterate previous proposition:

Proposition. Fix $k \in \mathbb{P}$ and $f_1, \dots, f_k : \mathbb{N} \rightarrow \mathbb{C}$. Define $h : \mathbb{N} \rightarrow \mathbb{C}$ by

$$h(\#X) = \sum f_1(\#S_1) \cdots f_k(\#S_k),$$

where $\cup S_i = X$ and $S_i \cap S_j = \emptyset$ if $i \neq j$. Then

$$E_h(x) = E_{f_1}(x) \cdots E_{f_k}(x).$$

A **partition** of a finite set S is a collection $\{B_1, \dots, B_k\}$ of subsets (called **blocks**) of S such that

$$\cup B_i = S, \quad B_i \neq \emptyset, \quad B_i \cap B_j = \emptyset \text{ if } i \neq j.$$

Write $\Pi(S)$ for the set of partitions of S .

Partitions of $[3]$:

$$1 - 2 - 3 \quad 12 - 3 \quad 13 - 2 \quad 1 - 23 \quad 123$$

Exponential formula. Given $f : \mathbb{P} \rightarrow \mathbb{C}$, define $h : \mathbb{N} \rightarrow \mathbb{C}$ by

$$h(0) = 1$$

$$h(\#S) = \sum_{\pi} f(\#B_1) \cdots f(\#B_k), \quad \#S > 0,$$

where $\pi = \{B_1, \dots, B_k\} \in \Pi(S)$. Then

$$E_h(x) = e^{E_f(x)}.$$

Proof. Set $f(0) = 0$. For **fixed** k let

$$g_k(\#S) = \sum_{(B_1, \dots, B_k)} f(\#B_1) \cdots f(\#B_k),$$

where $\{B_1, \dots, B_k\} \in \Pi(S)$. Thus

$$E_{g_k}(x) = E_f(x)^k.$$

Since $T_i \neq \emptyset$, all $k!$ orderings of T_1, \dots, T_k are distinct. Thus for fixed k , if

$$h_k(\#S) = \sum_{\{B_1, \dots, B_k\} \in \Pi(S)} f(\#B_1) \cdots f(\#B_k),$$

then $E_{h_k}(x) = \frac{1}{k!} E_{g_k}(x) = \frac{1}{k!} E_f(x)^k$.

Hence

$$\begin{aligned} E_h(x) &= 1 + \sum_{k \geq 1} E_{h_k}(x) \\ &= \sum \frac{E_f(x)^k}{k!} = e^{E_f(x)}. \quad \square \end{aligned}$$

Examples. (a) Let $\Pi_n = \Pi([n])$
and $B(n) = \#\Pi_n$ (**Bell number**).
If $f(i) = 1 \forall i$ then

$$B(n) = \sum_{\{B_1, \dots, B_k\} \in \Pi_n} f(\#B_1) \cdots f(\#B_k).$$

Thus

$$\begin{aligned} \sum_{n \geq 0} B(n) \frac{x^n}{n!} &= \exp \sum_{n \geq 1} \frac{x^n}{n!} \\ &= \exp(e^x - 1), \end{aligned}$$

(b) Let $f(n)$ be the number of **con-**
nected graphs on the vertex set $[n]$.
Thus $h(n)$ is the **total** number of graphs
on $[n]$, so $h(n) = 2^{\binom{n}{2}}$. Hence

$$\sum_{n \geq 1} f(n) \frac{x^n}{n!} = \log \sum_{n \geq 0} 2^{\binom{n}{2}} \frac{x^n}{n!}.$$


(Note that these series **diverge** for all
 $x \neq 0$.)

(c) Let $t_k(n)$ be the number of permutations w of $[n]$ satisfying $w^k = 1$. Thus every cycle length d of w satisfies $d|k$. We can choose w by partitioning $[n]$ into blocks of sizes $d|k$ and placing a cycle on each such block in $(d-1)!$ way. Hence


$$\begin{aligned} \sum_{n \geq 0} t_k(n) \frac{x^n}{n!} &= \exp \sum_{d|k} (d-1)! \frac{x^d}{d!} \\ &= \exp \sum_{d|k} \frac{x^d}{d}. \end{aligned}$$

TREES


A **rooted tree** is a connected graph without cycles with one distinguished vertex (the **root**).



Let $r(n)$ be the number of rooted trees on the vertex set $[n]$. E.g., $r(3) = 9$:



To obtain a rooted tree T on $[n]$, choose a root r in n ways, choose a partition $\pi \in \Pi([n] - \{r\})$, place a rooted tree T_i on each block of π , and “join” r to the roots of each T_i .



Let

$$\mathbf{R}(x) = \sum_{n \geq 1} r(n) \frac{x^n}{n!}$$
$$e^{\mathbf{R}(x)} = \sum_{n \geq 0} \mathbf{f}(n) \frac{x^n}{n!}.$$

Thus $f(n)$ is the number of **forests of rooted trees** on $[n]$, so $xe^{\mathbf{R}(x)}$ is the exponential generating function for choosing a 1-element subset of $[n]$ (the root) and placing a forest of rooted trees on the remaining elements. Since this structure is equivalent to a rooted tree on $[n]$, we have

$$\mathbf{R}(x) = xe^{\mathbf{R}(x)}.$$

Given

$$F(x) = a_1x + a_2x^2 + \cdots, \quad a_1 \neq 0,$$

define $F(x)^{\langle -1 \rangle}$ by


$$F(F^{\langle -1 \rangle}(x)) = F^{\langle -1 \rangle}(F(x)) = x$$

(exists and is unique). Then

$$\begin{aligned} R(x) &= xe^{R(x)} \\ \Rightarrow R(x) &= (xe^{-x})^{\langle -1 \rangle}. \end{aligned}$$


How to find the coefficients $r(n)/n!$ of $(xe^{-x})^{\langle -1 \rangle}$?

Bijective proof (Joyal). A **double rooted tree** is a tree with one vertex labelled s (start) and one vertex (possibly the same) labelled e (end). The number of double rooted trees on $[n]$ is $n \cdot r(n)$. Let T be such a tree, and let P be the unique path from s to e .




The vertices from s to e form a permutation of its elements written in increasing order. Write this permutation in cycle form as a directed graph:

1	3	6	9	11	12	16
12	6	16	9	3	1	11



Attach the subtrees of the path P back to their attached vertices and directed into the cycles:



We obtain a digraph on $[n]$ for which every vertex has outdegree one, i.e., the graph of a function $f : [n] \rightarrow [n]$. Conversely, every such f comes from a unique double rooted tree T .

$$\begin{aligned} & \# \text{ of } f : [n] \rightarrow [n]: \mathbf{n^n} \\ \Rightarrow & \# \text{ double-rooted trees on } [n]: \mathbf{n^n} \\ & \Rightarrow r(n) = \mathbf{n^{n-1}} \end{aligned}$$

Can we generalize this argument to find coefficients of other $F^{\langle -1 \rangle}(x)$?

Lagrange inversion formula. *Let*

$$F(x) = a_1x + a_2x^2 + \cdots, \quad a_1 \neq 0.$$

Let $k, n \in \mathbb{Z}$. Then

$$n[x^n]F^{\langle -1 \rangle}(x)^k = k[x^{n-k}] \left(\frac{x}{F(x)} \right)^n.$$

Proof. A combinatorial proof can be given based on counting trees. Proof of Lagrange:

Consider **Laurent series**

$$G(x) = \sum_{n \geq n_0 \in \mathbb{Z}} b_n x^n.$$

For instance,

$$\begin{aligned} \frac{1}{F(x)^k} &= \frac{1}{(a_1x + a_2x^2 + \dots)^k} \\ &= \frac{1}{x^k(a_1 + a_2x + \dots)^k} \\ &= x^{-k}(d_0 + d_1x \dots) \\ &= d_0x^{-k} + d_1x^{-k+1} + \dots \end{aligned}$$

Key fact:

$$[x^{-1}] \frac{d}{dx} G(x) = 0$$

$$\text{Set } F^{\langle -1 \rangle}(x)^k = \sum_{i \geq k} p_i x^i, \text{ so}$$

$$x^k = \sum_{i \geq k} p_i F(x)^i.$$

Apply $\frac{d}{dx}$:

$$kx^{k-1} = \sum_{i \geq k} ip_i F(x)^{i-1} F'(x)$$

$$\Rightarrow \frac{kx^{k-1}}{F(x)^n} = \sum_{i \geq k} ip_i F(x)^{i-n-1} F'(x).$$

Take $[x^{-1}]$ on both sides. Since

$$F(x)^{i-n-1}F'(x) = \frac{1}{i-n} \frac{d}{dx} F(x)^{i-n}, \quad i \neq n,$$

the coefficient of x^{-1} of the right-hand side is

$$\begin{aligned} [x^{-1}]np_n \frac{F'(x)}{F(x)} &= [x^{-1}]np_n \left(\frac{a_1 + 2a_2x + \cdots}{a_1x + a_2x^2 + \cdots} \right) \\ &= [x^{-1}]np_n \left(\frac{1}{x} + \cdots \right) \\ &= np_n. \end{aligned}$$

Hence

$$[x^{-1}] \frac{kx^{k-1}}{F(x)^n} = np_n = n[x^n]F^{\langle -1 \rangle}(x)^k,$$

which is equivalent to

$$n[x^n]F^{\langle -1 \rangle}(x)^k = k[x^{n-k}] \left(\frac{x}{F(x)} \right)^n. \quad \square$$

Let

$$R(x) = (xe^{-x})^{\langle -1 \rangle} = \sum_{n \geq 1} r(n) \frac{x^n}{n!}.$$

Thus if $r_k(n)$ is the number of forests of k rooted trees on $[n]$, then

$$\frac{1}{k!} R(x)^k = \sum_{n \geq k} r_k(n) \frac{x^n}{n!}.$$

By Lagrange inversion,

$$\begin{aligned}n[x^n]R(x)^k &= k[x^{n-k}] \left(\frac{x}{xe^{-x}} \right)^n \\ &= k[x^{n-k}]e^{nx} \\ &= \frac{kn^{n-k}}{(n-k)!},\end{aligned}$$

so

$$\begin{aligned}r_k(n) &= \frac{k}{n} \frac{n!}{(n-k)!k!} n^{n-k} \\ &= \binom{n-1}{k-1} n^{n-k}.\end{aligned}$$

ALGEBRAIC FUNCTIONS

A power series $F(x) = a_0 + a_1x + \dots$ is **algebraic** if \exists a polynomial $L(u, v) \neq 0$ such that

$$L(x, F(x)) = 0.$$

Examples. (a) Rational functions $F(x) = P(x)/Q(x)$ are algebraic, since

$$Q(x)F(x) - P(x) = 0.$$

(b) Easy to check that

$$\binom{-1/2}{n} = \left(-\frac{1}{4}\right)^n \binom{2n}{n},$$

so

$$F(x) := \sum_{n \geq 0} \binom{2n}{n} x^n = \frac{1}{\sqrt{1-4x}}.$$

(c) Let $F(x) = \sum_{n \geq 0} \binom{3n}{n} x^n$. Then

$$(27x - 4)F(x)^3 + 3F(x) + 1 = 0.$$

(d) **Not** algebraic:

$$\sum_{n \geq 0} \binom{2n}{n}^2 x^n, \quad \sum_{n \geq 0} \frac{(3n)!}{n!^3} x^n.$$

Theorem. Let $F(x) = \sum_{n \geq 0} f(n)x^n$ be algebraic. Then $\exists d \geq 1$ and polynomials $P_0(n), \dots, P_d(n)$ (not all 0) such that for all $n \geq 0$

$$P_d(n)f(n+d) + P_{d-1}(n)f(n+d-1) \\ + \dots + P_0(n)f(n) = 0.$$

One says $F(x)$ is **D-finite** and $f(n)$ is **P-recursive**.

Proof (sketch). Let $L(u, v)$ be a nonzero polynomial such that $L(x, F(x)) = 0$.

Thus

$$\begin{aligned} L_u(x, F(x)) + F'(x)L_v(x, F(x)) &= 0 \\ \Rightarrow F'(x) &= -\frac{L_u(x, F(x))}{L_v(x, F(x))} \in \mathbb{C}(x, F(x)). \end{aligned}$$

Similarly all higher derivatives $F^{(i)}(x) \in \mathbb{C}(x, F(x))$. Since $F(x)$ is algebraic

$$\dim_{\mathbb{C}(x)} \mathbb{C}(x, F(x)) < \infty.$$

Thus $F(x), F'(x), F''(x), \dots$ are linearly dependent over $\mathbb{C}(x)$. Write down this linear dependence relation, clear denominators, and equate coefficients of x^n to get an equation

$$P_d(n)f(n+d) + \dots + P_0(n)f(n) = 0. \quad \square$$

Example. Let $f(m, n)$ be the number of paths from $(0, 0)$ to (m, n) with steps $(1, 0)$, $(0, 1)$, $(1, 1)$ (**Delannoy number**). Thus

$$\begin{aligned} \sum_{m, n \geq 0} f(m, n) x^m y^n &= \sum_{k \geq 0} (x + y + xy)^k \\ &= \frac{1}{1 - x - y - xy}. \end{aligned}$$

Then

$$y := \sum_{n \geq 0} f(n, n)x^n = [t^0] \frac{1}{1 - xt - \frac{1}{t} - x}$$

$$= [t^0] \frac{1}{\beta - \alpha} \left(\frac{t}{t - \alpha} - \frac{t}{t - \beta} \right),$$

where $\alpha = \frac{1}{2}(1 - x - \sqrt{1 - 6x + x^2})$,
 $\beta = \frac{1}{2}(1 - x + \sqrt{1 - 6x + x^2})$. Hence

$$y = [t^0] \frac{1}{\sqrt{1 - 6x + x^2}} \left(\frac{t\alpha^{-1}}{1 - t\alpha^{-1}} + \frac{1}{1 - t^{-1}\beta} \right)$$

$$= \frac{1}{\sqrt{1 - 6x + x^2}},$$

and we get for $g(n) = f(n, n)$,

$$(n+2)g(n+2) - 3(2n+3)g(n+1) + (n+1)g(n) = 0$$

(challenging to prove directly!).

k -ARY PLANE TREES

A k -ary plane tree is a rooted tree for which every non-endpoint vertex has k cyclically ordered subtrees.

Let $f_k(n)$ denote the number of k -ary plane trees with n vertices and

$$y = F_k(x) = \sum_{n \geq 0} f_k(n)x^n.$$

Then $y = x + xy^k$, so

$$y = \left(\frac{x}{1 + x^k} \right)^{\langle -1 \rangle}.$$

By Lagrange inversion,

$$\begin{aligned} n[x^n]y &= [x^{n-1}](1 + x^k)^n \\ \Rightarrow f_k(n) &= \begin{cases} \frac{1}{n} \binom{n}{j}, & n = kj + 1 \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Special case: $k = 2$ (plane **binary** trees). Then

$$f_2(2n + 1) = \frac{1}{n + 1} \binom{2n}{n},$$

a **Catalan number** C_n .


66 combinatorial interpretations of C_n :
Exercise 6.19 of *Enumerative Combinatorics*, vol. 2.

36 additional interpretations (as of 22
December 2002):

www-math.mit.edu/~rstan/ec

Examples.


- triangulations of a convex $(n+2)$ -gon into n triangles by $n - 1$ diagonals that do not intersect in their interiors




- binary parenthesizations of a string of $n + 1$ letters

$(xx \cdot x)x$ $x(xx \cdot x)$ $(x \cdot xxx)x$ $x(x \cdot xxx)$ $xx \cdot xx$

- lattice paths from $(0, 0)$ to (n, n) with steps $(0, 1)$ or $(1, 0)$, never rising above the line $y = x$




- n nonintersecting chords joining $2n$ points on the circumference of a circle



- permutations $a_1 a_2 \cdots a_n$ of $[n]$ with longest decreasing subsequence of length at most two (i.e., there does not exist $i < j < k, a_i > a_j > a_k$)

123 213 132 312 231


- ways to stack coins in the plane, the bottom row consisting of n consecutive coins



- n -tuples (a_1, a_2, \dots, a_n) of integers $a_i \geq 2$ such that in the sequence $1a_1a_2 \cdots a_n1$, each a_i divides the sum of its two neighbors

14321 13521 13231 12531 12341

Bijjective proof that there are $C_n = \frac{1}{n+1} \binom{2n}{n}$ plane binary trees with $2n + 1$ vertices: do a depth-first (preorder) search through the tree, labeling down edges 1, up edges -1 , and ignoring the last edge.



1 1 - 1 1 - 1 - - - 1 1

This converts trees to sequences of $n + 1$ 1's and $n - 1$'s such that every partial sum is positive.

Claim. For any sequence $a_1 a_2 \cdots a_{2n+1}$ of $n + 1$ 1's and $n - 1$'s, there is exactly one value of i for which every partial sum of $a_i a_{i+1} \cdots a_{2n+1} a_1 a_2 \cdots a_{i-1}$ is positive.

Claim. For any sequence $a_1 a_2 \cdots a_{2n+1}$ of $n + 1$ 1's and $n - 1$'s, there is exactly one value of i for which every partial sum of $a_i a_{i+1} \cdots a_{2n+1} a_1 a_2 \cdots a_{i-1}$ is positive.

Proof (naive). Induction on n . Clear for $n = 0$. Assume for $n - 1$. Given $\alpha = a_1 \cdots a_{2n+1}$, can always find $a_j = 1$, $a_{j+1} = -1$ (subscripts modulo $2n + 1$). Remove a_j, a_{j+1} from α , giving $\beta = b_1 \cdots b_{2n-1}$. By the induction hypothesis there is a unique i for which $b_i \cdots b_{i-1}$ has all partial sums positive. If $b_i = a_k$, then k is the unique integer for which $a_k \cdots a_{k-1}$ has every partial sum positive. \square

There are $\binom{2n+1}{n}$ sequences of $n + 1$ 1's and $n - 1$'s. All their $2n + 1$ “cyclic shifts” are distinct since $\gcd(n, n + 1) = 1$. Thus the number of plane binary trees with $2n + 1$ vertices is

$$\frac{1}{2n + 1} \binom{2n + 1}{n} = \frac{1}{n + 1} \binom{2n}{n} = C_n.$$